www.coastaldigestivecare.com





Colonoscopy Preparation Planner

Procedure Time: you will be contacted within 1 week of the procedure date with your arrival time.

Transportat

<u>Transportation</u>: Once you have received your arrival time, coordinate your transportation needs for the day of the procedure. According to State law you may not, under any circumstance, be allowed to drive a motor vehicle home or walk home once your procedure is completed. We reserve the right to cancel the procedure at any time if transportation is not confirmed.



<u>Medications</u>: Please stop Vitamin E, any multi-vitamins containing Vitamin E, any oil based supplements, (fish oil, castor oil), and Iron supplements 1 week prior to procedure. <u>DIABETICS SHOULD CHECK WITH THEIR DOCTOR REGARDING THEIR MEDICATIONS.</u>



3 days before procedure:

Begin the recommended low residue diet. (See website for more information on low residue diets)



2 days before procedure: (Last Chance to cancel procedure without possible financial penalties)

- 1. **Begin the recommended clear liquids diet at 6 PM.** (See website for more information on clear liquids)
- 2. Drink a minimum of (8) glasses of CLEAR liquids (total of ~64oz.), throughout the day.
- 3. Eat your last meal between the hours of 5-6 PM.



1 Day before your procedure:

- 1. Continue the recommended clear liquid diet. (See website for details)
- 2. Drink a minimum of (8) glasses of CLEAR liquids (total of ~64oz.), throughout the day to prevent dehydration.



6:30-10:30 Arrival times:

FIRST STEP- 4PM (day before procedure), take (1) Ducolax® tablet with water.

SECOND STEP- 6PM (day before procedure) Begin drinking HalfLytely® solution. *Drink (1) 8oz.

glass every 10 minutes over 1 hour (~ 8 glasses).

10:45-3:30 Arrival times:

FIRST STEP- 4PM (day before procedure) Take (1) Ducolax® tablet with water.

SECOND STEP- 6PM (day before procedure) Begin drinking HalfLytely® solution. *Drink (1) 8oz.

glass every 10 minutes over 1 ½ hours (~4 glasses).

THIRD STEP- 6AM (day of procedure) Begin drinking the remaining HalfLytely® solution.

*Drink (1) 8oz. glass every 10 minutes over 40 minutes (~4 glasses).

** Apply Vaseline or hemorrhoid cream/wipes to any anal irritation that may occur.**

Procedure Day:

- 1. If your bowel movements are not clear take 1-2 Fleet® enemas before leaving home. (Clear can be defined as yellow or water-clear).
- 2. Arrive at your appointed time wearing loose-fitting clothing, socks, and a short-sleeved T-shirt.



3. NOTHING TO DRINK WITHIN 4 HOURS OF YOUR ARRIVAL TIME.